


# The Acene Series: Narrowest Zig-Zag Graphene Nanoribbons

Prof. Dr. Holger F. Bettinger

Institut für Organische Chemie, Auf der Morgenstelle 18, 72076 Tübingen, Germany

Acenes [n]ac, consisting of linearly fused benzene rings, are important materials for organic electronics and are currently in the focus of materials research.<sup>[1]</sup> Reactivity increases among the acene series significantly with length,<sup>[2]</sup> causing instability for acenes longer than hexacene.



The acenes beyond hexacene are assumed to be notoriously unstable,<sup>[3]</sup> but are valuable for studying the transition from stable organic molecules to zig-zag graphene nanoribbons that may be metallic in the limit of infinite length. To allow experimental investigations of larger acenes,<sup>[4]</sup> we have developed synthetic access to acenes up to undecacene [11]ac.<sup>[5]</sup> We report on the properties of those higher acenes, in particular with respect to their optical and magnetic properties.

## References

- [1] a) J. E. Anthony, *Angew. Chem. Int. Ed.* **2008**, *47*, 452-483; b) J. E. Anthony, *Chem. Rev.* **2006**, *106*, 5028-5048; c) F. Ortmann, K. S. Radke, A. Günther, D. Kasemann, K. Leo, G. Cuniberti, *Adv. Funct. Mater.* **2015**, *25*, 1933-1954.
- [2] D. Biermann, W. Schmidt, *J. Am. Chem. Soc.* **1980**, *102*, 3163-3173.
- [3] a) R. Einholz, H. F. Bettinger, *Angew. Chem., Int. Ed.* **2013**, *52*, 9818-9820; b) R. Einholz, T. Fang, R. Berger, P. Grüninger, A. Früh, T. Chassé, R. F. Fink, H. F. Bettinger, *J. Am. Chem. Soc.* **2017**, *139*, 4435-4442.
- [4] H. F. Bettinger, C. Tönshoff, *Chem. Rec.* **2015**, *15*, 364-369.
- [5] a) C. Tönshoff, H. F. Bettinger, *Angew. Chem., Int. Ed.* **2010**, *49*, 4125-4128; b) B. Shen, J. Tatchen, E. Sanchez-Garcia, H. F. Bettinger, *Angew. Chem. Int. Ed.* **2018**, *57*, 10506-10509.